

# Index

The index does not include the data provided in the annexes to the chapters.

Boxes, figures, and tables are indicated by b, f, and t after the page numbers.

abortion, non-use or ineffective use of  
    contraception as risk factor for, 391t  
addiction. *See* alcohol use; drug abuse;  
    smoking  
Africa. *See* Sub-Saharan Africa  
age  
    burden of disease and age differences,  
        88–89  
    deaths by cause, region, sex, and age,  
        126–179t  
    regional distribution of population  
        by, 19–20, 22f  
risk factors and burden of  
    disease, 247, 251  
PAFs, deaths, YLL, and DALYs,  
    269–393t  
weighting  
    in DALY measure, 400–401, 401f, 423  
    in YLL, 401–402, 401f, 402f, 402t  
AIDS. *See* HIV/AIDS  
air pollution as risk factor, 268, 372–375t  
    indoor smoke as risk factor, 362–365t  
alcohol use  
    as cause of burden of disease, 9, 85, 247  
    child sexual abuse as risk factor, 356t  
    code to describe death due to, 58  
    counterfactual analysis of, 246  
    incidence and prevalence, data  
        sources on, 82  
        as risk factor, 325–347t  
Alzheimer's disease, data sources on  
    incidence and prevalence, 83  
anemia  
    incidence and prevalence, data  
        sources on, 81  
        as risk factor, 247, 276–279t  
angina. *See* cardiovascular disease (CVD)  
anxiety disorders. *See* neuropsychiatric  
    disorders  
*Armed Conflict Report* on war deaths by  
    country and year, 65

arthritis  
    incidence and prevalence, data sources  
        on, 84–85  
    obesity as risk factor, 306t  
Asia. *See specific countries and regions*  
asthma, data sources on incidence and  
    prevalence, 84  
avian influenza threat, 162  
biases in data and uncertainty, 408  
bladder cancer, smoking as risk factor, 380t  
blindness. *See* vision loss and blindness  
blood pressure. *See* cardiovascular  
    disease (CVD)  
breast cancer  
    alcohol use as risk factor, 329t  
    as leading cause of cancer death for  
        women, 90  
    obesity as risk factor, 300t  
    physical inactivity as risk factor, 316t  
burden of disease  
    *See also* DALYs (disability-adjusted life  
        years); Global Burden of Disease  
        (GBD) framework; mortality rates;  
        years of healthy life lost as result of  
        disability (YLD); years of life lost  
        (YLL); *specific diseases and conditions*  
    analyses as information tools for health  
        policy, 4–5  
    by cause, 7–10, 8t, 88, 88t, 108–118t  
    causes and codes, 56–59, 57f, 58t, 95–99t  
    history of, 2–4  
    income levels and, 89f, 89t  
    leading causes of, 85–86, 86t, 91t  
    discounting and age weighting. *See* age;  
        discounting  
    risk factors, 247, 251–252, 251f, 255t  
    risk factors for, 5, 242, 247–252, 248f,  
        249–250t  
    sensitivity and uncertainty analyses,  
        399–426

*See also* sensitivity analysis; uncertainty  
    analysis

cancer  
    air pollution as risk factor, 373t  
    alcohol use as risk factor, 326–330t  
    as cause of burden of disease, 89  
    contaminated injections in health care  
        setting as risk factor, 369t  
    data sources on  
        deaths due to, 64  
        incidence and prevalence, 81  
    disability weights for, 124t  
    ill-defined codes for death caused  
        by, 58–59  
    indoor smoke as risk factor, 363t  
    low fruit and vegetable intake as risk  
        factor, 308–311t  
    obesity as risk factor, 299–301t  
    physical inactivity as risk factor, 315–316t  
    risk factor clusters and, 259, 260–262t, 267  
    smoking as risk factor, 377–384t  
    unsafe sex as risk factor, 323t  
cardiovascular disease (CVD)  
    alcohol use as risk factor, 335–336t  
    as cause of death, 69  
    ill-defined codes for, 56–58, 57f, 58t  
        by region, 73  
    cholesterol as risk factor, 296t  
    disease burden of, 9, 89  
    high blood pressure as risk factor,  
        291–295t  
    incidence and prevalence, data sources  
        on, 83–84  
    joint hazards of risk factors for, 253–254  
    low fruit and vegetable intake as risk  
        factor, 312t  
    obesity as risk factor, 303–304t  
    physical inactivity as risk factor, 318t  
    risk factor clusters and, 259, 263–264t  
    smoking as risk factor, 385–386t

- Caribbean. *See* Latin America and Caribbean
- categorical attribution, 242
- cause of death models, 51
- for countries with poor data, 66–68
- Central Asia. *See* Europe and Central Asia
- cerebrovascular disease
- alcohol use as risk factor, 337t
  - as cause of death and disease burden, 89, 90
  - cholesterol as risk factor, 297t
  - high blood pressure as risk factor, 293t
  - incidence and prevalence, data sources on, 84
  - as leading cause of death, 72
  - low fruit and vegetable intake as risk factor, 313t
  - obesity as risk factor, 305t
  - physical inactivity as risk factor, 319t
  - smoking as risk factor, 387t
- cervical cancer
- smoking as risk factor, 379t
  - unsafe sex as risk factor, 323t
- Chagas disease, data on deaths due to, 64
- CHERG. *See* Child Health Epidemiology Reference Group
- childbirth conditions
- acquisition of life potential (ALP), 431, 438–439, 439t
  - burden of disease resulting from, 431–441
  - death rates due to, 54
  - low birthweight
    - alcohol use as risk factor, 325t
    - preterm birth and, 431
  - maternal death, 64
  - neonatal death and stillbirth, 427–463
    - See also* child mortality rates
- Child Health Epidemiology Reference Group (CHERG; WHO), 34, 424, 430–431, 461
- childhood health
- See also* child mortality rates; specific diseases
  - environmental risk factors and child undernutrition, 254
  - hearing loss, 85
  - malnutrition, 33, 247
  - underweight as risk factor, 269–275t
- child mortality rates
- acute respiratory infections as cause, 63
  - communicable diseases and, 68
  - data available on, 21–22, 24, 32, 35, 53, 64
  - diarrheal disease and, 31
  - HIV/AIDS and, 26, 31, 33, 71
  - injuries as cause of death, 32, 34
  - leading causes of child deaths, 69–71
    - under age five, 28, 29–30t, 31f
    - trends in, 28–32, 35
  - lowering of, 26
  - malaria and, 31
  - neonatal mortality and stillbirths, 7, 427–463
    - cause of deaths, 430–431, 445–460t
    - cause of stillbirths, 431
    - childbirth conditions and, 431–441
    - comparison of GBD and CHERG numbers, 461
    - in context of global burden of disease, 428–431, 441t, 448t
    - DALYs and, 431, 438–441, 438f, 439f, 441t, 442–444, 452–454t
    - deaths by age and cause, 432–437t
    - number of deaths and death rates, 428–430, 429f, 429t, 430t
    - preterm birth and, 431
    - terminology, 428
    - YLL and, 440f, 440t
  - perinatal causes, 32, 33–34, 64, 69, 71
  - risk factor clusters and, 259, 265–266t, 267
  - vaccine-preventable diseases as cause, 31, 63
  - child sexual abuse as risk factor, 247, 355–361t
  - China
    - death registration and reporting in, 51, 53, 59–60, 59f
    - neonatal deaths in, 33, 34
    - YLL and YLD rates in, 87  - cholesterol as risk factor, 296–298t
  - cirrhosis of liver
    - alcohol use as risk factor, 338t
    - contaminated injections in health care setting as risk factor, 370t
  - clean drinking water. *See* water, unsafe
  - CodMod software, 51, 66–68
  - colon and rectal cancer
    - low fruit and vegetable intake as risk factor, 310t
    - obesity as risk factor, 299t
    - physical inactivity as risk factor, 315t
  - communicable diseases
    - See also* specific diseases
    - as cause of burden of disease, 8t, 90
    - childhood underweight as risk factor, 274t
  - data sources and methods for incidence and prevalence, 79–81
  - death rates due to, 53–54
  - risk factors for, 268
  - vitamin A deficiency as risk factor, 283t
- Comparative Risk Assessment (CRA) project, 241–247
- See also* risk factors
- counterfactual analysis used in, 242
- exposure and hazard, data on, 245–246
- framework, 242
- mortality estimates and burden of disease, 245
- population attributable fractions (PAFs) and estimating, 245
- by region, risk factor, disease outcome, age, and sex, 269–393t
- risk correlation, 254
- risk factor selection, 242–247
- congestive heart failure. *See* cardiovascular disease (CVD)
- contaminated injections in health care setting as risk factor, 366–371t
- contraception, non-use or ineffective use as risk factor, 247, 391–393t
- counterfactual analysis, 242
- CVD. *See* cardiovascular disease
- DALYs (disability-adjusted life years), 3b, 47–49
- age weighting in, 400–401, 401f, 423
- variations in parameter values, 402–406, 402f, 403t, 404f, 405f, 405t, 406f
- by cause, 88, 88t
- by cause, region, sex, and age, 180–233t
- childbirth conditions and, 431, 438–439, 438f, 439f
- comparison of high-income to low- and middle-income countries, 87f, 88f, 89f
- defined, 3b, 45
- discounting in, 400, 423
- variations in parameter values, 402–406, 403t, 404f, 405f, 405t, 406f
- by region, 87–90, 87f, 88f, 91t
- risk factors and burden of disease, 269–393t
- variations in parameter values, 408f
- social value choices and, 49–50
- stillbirths and, 442–444, 458–460t
- uncertainty ranges for disability weights, 413, 416–417, 416f, 417f, 418–419t, 420f, 421t
- use of, 3b, 45, 47–49
- data sources. *See* sources of data
- death. *See* mortality rates
- debate of appropriate values and priorities for burden of disease, 4
- dehydration. *See* diarrheal diseases

Demographic and Health Survey (DHS) program, 51

demographic trends of major regions (1990–2001), 18–21, 35, 36–42t

*See also* mortality rates

distribution by age and location, 19–20, 22f

fertility, 20–21, 20–21t

percentage of regional population covered by censuses, 18t

population size and growth, 19, 20t

sex ratios, 20, 23f

sources of data and methodology, 18–19

urbanization, 20

dengue, data sources on incidence and prevalence, 80

depression

- alcohol use as risk factor, 332t
- child sexual abuse as risk factor, 355t
- data sources for incidence and prevalence, 82
- as leading cause of disability, 7, 85

diabetes

- alcohol use as risk factor, 331t
- as cause of death, 69
- incidence and prevalence, data sources on, 81–82
- increasing burden of, 9, 90
- obesity as risk factor, 302t
- physical inactivity as risk factor, 317t

diarrheal diseases

- childhood underweight as risk factor, 269t
- child deaths due to, 31, 33
- data sources on

  - deaths due to, 33, 63
  - incidence and prevalence, 79

- deaths due to, 71, 247
- unsafe water, sanitation, and hygiene as risk factors, 353t
- vitamin A deficiency as risk factor, 280t
- zinc deficiency as risk factor, 287t

diphtheria, data on deaths due to, 63

disability

- See also* DALYs (disability-adjusted life years); years of healthy life lost as result of disability (YLD)
- defined, 51
- discounting

  - in DALY measure, 400, 423
  - variations in parameter values, 402–406, 403t, 404f, 405f, 405t, 406f
  - in YLL measure, 401–402, 401f, 402t

Disease Control Priorities Project, regional reporting categories for, 94t

DisMod, use to ensure consistency in YLD calculations, 74–75, 75f

drowning, alcohol use as risk factor, 342t

drug abuse

- child sexual abuse as risk factor, 357t
- data sources on

  - deaths due to, 64–65
  - incidence and prevalence, 82
  - as risk factor, 348–352t

drug resistance, 32

East Asia and Pacific

- causes of burden of disease in, 90
- DALYs by cause, sex, and age, 186–191t
- deaths by cause, sex, and age, 132–137t
- mortality rates in, 28
- pulmonary disease in, 73
- YLL and YLD rates in, 87
- economic growth and health, 26–27
- edentulism, data sources on incidence and prevalence, 85
- Egypt, death registration and reporting in, 60
- Ellison Institute for Global Health, 93
- environmental risk factors and child undernutrition, 254
- epidemiological assessments of cause of death, 51
- epidemiological studies as data source

  - analysis categories by region, 107t
  - lack of, 92
  - for YLD calculations, 76

- epilepsy

  - alcohol use as risk factor, 333t
  - incidence and prevalence, data sources on, 82–83

- esophageal cancer

  - alcohol use as risk factor, 327t
  - low fruit and vegetable intake as risk factor, 308t

- estimates. *See* mortality rates; sensitivity analysis; uncertainty analysis
- Europe and Central Asia

  - See also* specific countries
  - CVD in, 73
  - DALYs by cause, sex, and age, 192–197t
  - deaths by cause, sex, and age, 138–143t
  - demographic structure by age, 19
  - disease burden in, 9, 91
  - injuries as cause of burden of disease in, 91
  - mortality rates, trends in, 26, 27
  - sex ratio in, 20, 23f
  - YLL and YLD rates in, 87

fall-related injuries. *See* injuries

fertility trends, 20–21, 20–21t

gender differences

- age weighting and discounting for YLL, 402f
- birth ratio of boys to girls, 20
- burden of disease and, 88–89
- causes of death, 69, 70t, 126–179t
- injuries as cause of health burden, 90, 91–92
- mortality rates, 26–27t
- uncertainty estimates and, 410, 410f
- neuropsychiatric disorders, 85
- risk factors and burden of disease for PAFs, deaths, YLL, and DALYs, 269–393t
- traffic accidents, 90, 91

*Global Burden of Disease and Risk Factors (GBD). See* World Bank

Global Burden of Disease (GBD) framework, 1–2, 2f, 46–47

Handicap International's annual report on landmine victims, 65

health expectancies and health gaps, 47–48, 47f

health facility data as data source for YLD calculations, 76, 79

health status

- how to describe, 50–51
- population change and, 17

hearing loss

- incidence and prevalence, data sources on, 83
- as leading cause of YLD, 85, 92

heart disease. *See* cardiovascular disease (CVD)

helminth infections, data sources on

- incidence and prevalence, 80

hepatitis B

- contaminated injections in health care setting as risk factor, 367t
- incidence and prevalence, data sources on, 79

hepatitis C

- contaminated injections in health care setting as risk factor, 368t
- incidence and prevalence, data sources on, 79

high-income countries' burden of disease comparison to low- and middle-income countries. *See* DALYs (disability-adjusted life years); years of healthy life lost as result of disability (YLD)

DALYs by cause, sex, and age, 222–227t

- high-income countries' burden of disease (*Continued*)
- deaths by broad cause group, 68f, 69f
  - deaths by cause, sex, and age, 168–173t
  - leading causes of, 89f, 89t
  - risk factors by disease type, 251, 251f, 252
- HIV/AIDS
- See also specific countries and regions*
  - as cause of burden of disease, 90
  - child deaths due to, 26, 31, 33, 71
  - contaminated injections in health care setting as risk factor, 366t
  - deaths due to
    - data on, 63
    - in low- and middle-income countries, 7, 8, 68–69, 71  - drug abuse as risk factor, 348t
  - incidence and prevalence, data sources for, 79
  - unsafe sex as risk factor, 247, 252, 322t
- India
- death registration and reporting in, 51, 60
  - neonatal deaths in, 33
  - indoor smoke as risk factor, 362–365t
  - infant mortality. *See child mortality rates*
  - infectious diseases. *See communicable diseases*
  - information and surveillance
    - for knowledge to implement health policy, 4–5
    - for priority setting, 35  - injections in health care setting as risk factor, 366–371t
  - injuries
    - alcohol use as risk factor, 341t, 343–344t, 346t
    - as cause of health burden, 3, 7–9, 8t, 89, 90
    - child deaths due to, 32, 34
    - child sexual abuse as risk factor, 360t
    - data sources for incidence and prevalence, 85
    - disability weights for, 125t
    - drug abuse as risk factor, 350–351t
    - intent and codes assigned to death due to, 59
    - insomnia, data sources on incidence and prevalence, 82
    - Institute of Medicine (U.S.)
      - on quality of medical care, 166
      - on vaccine development priorities, 440    - International Campaign to Ban Landmines, 65
    - interventions, national priorities of, 4
    - intestinal nematode infections, data sources on incidence and prevalence, 80
    - iodine deficiency, data sources on incidence and prevalence, 81
    - Iran, death registration including cause of death, 5, 61, 61f
    - iron deficiency, 247
      - See also anemia*    - knowledge to implement health policy, 4–5
    - Landmine Monitor Report* on deaths due to landmines by country, 65
    - Latin America and Caribbean
      - See also specific countries*
      - DALYs by cause, sex, and age, 198–203t
      - deaths by cause, sex, and age, 144–149t
      - diabetes and endocrine disorders in, 9
      - HIV/AIDS in, 72
      - injuries as cause of burden of disease in, 91
      - mortality rates in, 28
      - violence in, 9, 73    - leprosy, data sources on incidence and prevalence, 80
    - leukemia, smoking as risk factor, 384t
    - life expectancy
      - See also years of life lost (YLL)*
      - age weighted and discounted YLL and, 402t
      - uncertainty estimates and, 409–411, 411f
      - YLD and, 86–87, 87f    - liver cancer
      - alcohol use as risk factor, 328t
      - contaminated injections in health care setting as risk factor, 369t
      - as leading cause of cancer deaths, 89–90
      - smoking as risk factor, 378t    - low- and middle-income countries' burden of disease
      - See also specific regions*
      - comparison to high-income countries.
        - See DALYs (disability-adjusted life years); years of healthy life lost as result of disability (YLD)*      - DALYs by cause, sex, and age, 180–185t
      - deaths by broad cause group, 68f, 69f
      - deaths by cause, sex, and age, 126–131t
      - epidemiological transition and reduction in, 90
      - leading causes of, 72t, 89f, 89t
      - neonatal mortality and stillbirths, 441t
      - noncommunicable diseases as cause of death, 7
      - risk factors by disease type, 251–252, 251f
      - lung cancer
        - air pollution as risk factor, 373t
        - indoor smoke as risk factor, 363t
        - as leading cause of cancer deaths, 69, 89      - low fruit and vegetable intake as risk factor, 311t
      - smoking as risk factor, 377t
      - lymphatic filariasis, data sources on incidence and prevalence, 80
      - malaria
        - as cause of burden of disease, 90
        - childhood underweight as risk factor, 271t
        - deaths due to
          - child deaths, 31
          - data on, 32–33, 64
          - in low- and middle-income countries, 8, 71        - incidence and prevalence, data sources for, 79
        - vitamin A deficiency as risk factor, 282t
        - zinc deficiency as risk factor, 288t      - males. *See gender differences*
      - malignant neoplasms. *See cancer*
      - malnutrition
        - childhood underweight as risk factor, 273t
        - decline in, 33
        - incidence and prevalence, data sources on, 81
        - joint hazards of child undernutrition for infectious diseases, 254
        - as leading risk for burden of disease, 247
        - as risk factor when low fruit and vegetable intake, 308–314t      - malpractice, 167
      - Mapping Malaria Risks in Africa International Collaboration, 64
      - maternal conditions
        - See also childbirth conditions*
        - anemia as risk factor, 276t
        - contraceptives, non-use or ineffective use as risk factor, 392t
        - incidence and prevalence, data sources on, 81
        - vitamin A deficiency as risk factor, 284t      - maternal mortality, data on death rate, 64
      - measles
        - childhood underweight as risk factor, 270t
        - child deaths due to, 31, 33
        - deaths due to, 71
          - data on, 33, 63
          - malnutrition and, 33        - vitamin A deficiency as risk factor, 281t      - Medical Outcomes Study (MOS) Short Form 36, 50
      - meningitis, data sources on incidence and prevalence, 79
      - mental retardation, data sources on incidence and prevalence, 83

Middle East and North Africa  
 DALYs by cause, sex, and age, 204–209  
 deaths by cause, sex, and age, 150–155t  
 fertility trends, 20  
 injuries as cause of burden of disease in, 91  
 mortality rates in, 28  
 traffic injuries in, 73, 90, 91  
 migraine, data sources on incidence and prevalence, 83  
 modeling to estimate cause of death for countries with poor data, 66–68  
 Monte Carlo simulation techniques, use in cause of death modeling, 67  
 mortality rates  
*See also* years of life lost (YLL); specific disease or condition as cause  
 cause of death models, 51  
 for countries with poor data, 66–68  
 causes of death, 7, 69–72, 70t  
 for adults, 71–72, 71t  
 for children, 28–32, 69, 71  
 in low- and middle-income countries, 126–131t  
 premature death, 73, 73f  
 by region, 72–73, 72t  
 changes in (1990–2001), 21–28, 32–34  
 characteristics by sex and region, 26, 26t  
 children. *See* child mortality rates  
 death registration information, 5, 22, 51, 52, 53, 55t, 92  
 distribution by major cause group, 68–69, 68f, 69f  
 epidemiological assessments of cause of, 51  
 estimates of, 19, 21–26, 424  
 differences of authors with UN data, 23–26, 24–25f  
 estimating by cause of death, 51–68  
 all-cause mortality data, 51–53, 52t  
 availability of data from countries, 21, 54–56, 55t  
 classification of causes of disease and injury, 53–54  
 countries with poor data, modeling for, 66–68  
 data sources and methods for specific countries, 59–61, 92, 100–106t  
 epidemiological estimates for specific causes, 61–66, 62t  
 ill-defined causes and “garbage codes,” 56–59, 57f, 57t, 58t  
 risk factors and attributable mortality, 245, 248f, 249–250t  
 traffic accidents as cause of, 69, 71, 73  
 trends in, 26–28, 35  
 true cause not reported, 54

uncertainty estimates and for all-cause mortality, 409–411, 410f, 411f  
 for regions by cause, 411–413, 413f, 414–415t  
 MOS (Medical Outcomes Study) Short Form 36, 50  
 mouth cancer, alcohol use as risk factor, 326t  
 Multiple Indicator Cluster Survey program, 51  
 multiple sclerosis, data sources on incidence and prevalence, 82–83  
 National Cancer Institute’s Surveillance, Epidemiology, and End Results program, 64  
 neonatal conditions and deaths. *See* childbirth conditions; child mortality rates  
 neuropsychiatric disorders as cause of health burden, 3–4, 7, 85, 90  
 data sources for incidence and prevalence, 83  
 newborn deaths. *See* child mortality rates  
 Nigeria, neonatal deaths in, 33  
 noncommunicable diseases  
*See also* specific diseases as cause of health burden, 3–4, 8, 8t, 9, 89–90, 91  
 incidence and prevalence, data sources on, 81–85  
 risk factors for, 268  
 obesity as risk factor, 247, 299–307t  
 onchocerciasis, data sources on incidence and prevalence, 80  
 osteoarthritis incidence and prevalence, data sources on, 85  
 obesity as risk factor, 306t  
 Pacific region. *See* East Asia and Pacific PAFs. *See* population attributable fractions  
 Pakistan, neonatal deaths in, 33  
 pancreas cancer, smoking as risk factor, 381t  
 panic disorder, child sexual abuse as risk factor, 359t  
 Parkinson’s disease, data sources on incidence and prevalence, 83  
 performance assessment and burden of disease as indicator, 4  
 perinatal conditions anemia as risk factor, 277t  
 incidence and prevalence, data sources on, 81  
 pertussis, data on deaths due to, 63  
 physical inactivity as risk factor, 315–320t  
 pneumonia, 72, 430–431  
 poisonings, alcohol use as risk factor, 340t  
 polio, data on deaths due to, 63  
 population attributable fractions (PAFs), 245, 269–393t  
 uncertainty and, 422–423, 422f  
 population surveys as data source for YLD calculations, 76  
 population trends. *See* demographic trends of major regions (1990–2001)  
 post-traumatic stress disorder, child sexual abuse as risk factor, 358t  
 premature death, 73, 73f  
 pulmonary disease, 73  
 air pollution as risk factor, 374t  
 incidence and prevalence, data sources on, 84  
 indoor smoke as risk factor, 364t  
 smoking as risk factor, 376t  
 quality-adjusted life year, defined, 47  
 rectal cancer. *See* colon and rectal cancer  
 registration of deaths. *See* mortality rates  
 resource allocation to generate greatest reduction in health loss, 5  
 respiratory infections air pollution as risk factor, 372t  
 childhood underweight as risk factor, 272t  
 child deaths due to, 31, 33  
 data sources on deaths, 63  
 incidence and prevalence, 80  
 indoor smoke as risk factor, 362t  
 smoking as risk factor, 388t  
 zinc deficiency as risk factor, 289t  
 rheumatoid arthritis, data sources on incidence and prevalence, 84–85  
 risk factors, 241–396  
*See also* Comparative Risk Assessment (CRA) project; specific factors (e.g., alcohol use, malnutrition)  
 adult vs. childhood diseases, 268  
 air pollution as, 372–375t  
 alcohol use as, 325–347t  
 anemia as, 276–279t  
 as cause of burden of disease, 4, 5, 9–10, 10t, 242, 247–252, 248f, 249–250t  
 expanding analysis to include factors not among leading causes, 268  
 joint effects of multiple factors, 255–259, 256–258t  
 ordering of, 247, 251–252, 251f, 255t  
 as cause of death, 248f, 249–250t  
 childhood underweight as, 269–275t  
 child sexual abuse as, 355–361t

- risk factors (*Continued*)
- cholesterol as, 296–298t
  - clusters, 259–267
  - contraceptives, non-use or ineffective use as, 391–393t
  - correlation, 254
  - data sources for mediated effects and effect modification, 253–254
  - drug abuse as, 348–352t
  - future research needs, 267
  - GBD framework assessing, 6
  - high blood pressure as, 291–295t
  - indoor smoke as, 362–365t
  - injections in health care setting as, 366–371t
  - joint effects of multiple factors, 252–267, 253f
  - low fruit and vegetable intake as, 308–314t
  - obesity as, 247, 299–307t
  - physical inactivity as, 315–320t
  - risk assessment, defined, 242
  - sanitation and hygiene as, 353–354t
  - sensitivity and uncertainty analyses, 406–407, 407f
  - smoking as, 376–390t
  - theoretical-minimum-risk exposure distributions, 242, 243–244t, 246–247
  - uncertainty and, 420–423
  - unsafe water as, 247, 353–354t
  - vitamin A deficiency as, 280–286t
  - zinc deficiency as, 287–290t
  - road accidents. *See* traffic injuries
  - Russian mortality rate, 26, 27, 87
  - sanitation and hygiene as risk factor, 353–354t
  - schistosomiasis, data sources on incidence and prevalence, 79–80
  - schizophrenia, data sources on incidence and prevalence, 82
  - sensitivity analysis, 399–407
  - age weighting
    - in DALY measure, 400–401, 401f, 423
    - variations in parameter values, 402–406, 402f, 403t, 404f, 405f, 405t, 406f
    - in YLL measure, 401–402, 401f, 402f  - discounting
    - in DALY measure, 400, 423
    - variations in parameter values, 402–406, 403t, 404f, 405f, 405t, 406f
    - in YLL measure, 401–402, 401f  - risk factors, 406–407, 407f
  - variations in parameter values, 408f
  - sepsis and neonatal deaths, 430–431
  - sexual abuse. *See* child sexual abuse as risk factor
  - sexually transmitted infections
    - See also* HIV/AIDS
    - incidence and prevalence, data sources on, 79
    - unsafe sex as risk factor, 247, 252, 321–324t  - smoking
    - burden of disease attributable to, 247
    - joint hazards with other risk factors, 254
    - as risk factor, 376–390t  - sources of data
    - criticism of GBD study based on, 423
    - death by cause, 51–68
    - demographic trends of major regions (1990–2001), 18–19
    - lack of, 92–93
    - uncertainty and, 408–409
    - years of healthy life lost as result of disability (YLD), 75–79  - South Asia
    - See also specific countries*
    - communicable diseases in, 90
    - DALYs by cause, sex, and age, 210–215t
    - deaths by cause, sex, and age, 156–161t
    - HIV/AIDS in, 72
    - noncommunicable diseases in, 90
    - sex ratio in, 20, 23f
    - suicide in, 73
    - YLL and YLD rates in, 87  - statistical methods and data, use in cause of death modeling, 66–67
  - stillbirths. *See* child mortality rates
  - stomach cancer
    - as leading cause of cancer deaths, 89
    - low fruit and vegetable intake as risk factor, 309t
    - smoking as risk factor, 382t  - strategic health planning, role of, 1
  - stroke. *See* cerebrovascular disease
  - Sub-Saharan Africa
    - See also specific countries*
    - child mortality rates in, 26, 32
    - communicable diseases in, 72
    - DALYs by cause, sex, and age, 216–221t
    - data sources on, 92
    - deaths by cause, sex, and age, 162–167t
    - fertility trends in, 20
    - HIV/AIDS in, 8, 26, 33, 71, 72, 90
    - malaria in, 8, 31, 72, 90
    - malnutrition in, 247
    - mortality rates in, 24–25, 26, 27
    - traffic accidents in, 90  - unsafe sex in, 247, 252
  - YLL and YLD rates in, 87
  - suicide, 73, 90, 91
  - surveillance. *See* information and surveillance
  - TB. *See* tuberculosis
  - tetanus and neonatal deaths, 431
  - Thailand
    - death registration and reporting in, 61
    - verbal autopsy study in, 5  - theoretical-minimum-risk exposure distributions. *See* risk factors
  - tobacco use. *See* smoking
  - trachoma, data sources on incidence and prevalence, 80
  - traffic injuries
    - alcohol use as risk factor, 339t
    - as cause of death, 9, 69, 71, 73
    - as cause of health burden, 90, 91  - tuberculosis (TB), data sources on deaths, 62–63
    - incidence and prevalence, 79  - Turkey, death registration and reporting in, 60–61
  - uncertainty analysis, 399, 408–409
    - criticism of GBD study based on, 423
    - cross-national comparisons and, 7, 54
    - describing and quantifying uncertainty, 409  - disability weights and, 413, 416–417, 416f, 417f, 418–419t, 420f, 421t
  - estimates
    - for all-cause mortality and life expectancies, 409–411, 410f, 411f
    - from epidemiological estimates, 417, 420
    - for regions by cause, 411–413, 413f, 414–415t  - PAFs and, 422–423, 422f
  - risk factors and, 420–423
  - sources of uncertainty, 408–409
  - undernutrition. *See* malnutrition
  - United Nations Population Division, 18–19, 23–26, 24–25f
  - United Nations Programme on HIV/AIDS, 63
  - unsafe sex. *See* sexually transmitted infections
  - unsafe water as risk factor, 247, 353–354t
  - urbanization trends, 20
  - uterine cancer, obesity as risk factor, 301t
  - vaccine-preventable diseases
    - See also specific disease*

data on deaths due to, 63  
incidence and prevalence, data sources on, 79  
verbal autopsies, 5, 28, 32, 92  
violence  
  alcohol use as risk factor, 345t  
  as cause of death, 71, 73  
  as cause of health burden, 9, 90, 91  
vision loss and blindness  
  incidence and prevalence, data sources on, 83  
  as leading cause of YLD, 85  
vitamin A deficiency  
  incidence and prevalence, data sources on, 81  
  as risk factor, 280–286t  
  
war, deaths due to, 56, 65–66  
water, unsafe, as risk factor, 247, 353–354t  
women  
  *See also* childbirth conditions; gender differences; maternal conditions  
breast cancer, 90  
death in childbirth, 64  
depression and, 7  
injuries as cause of death of, 89, 91–92  
world  
  DALYs by cause, sex, and age, 228–233t  
  deaths by cause, sex, and age, 174–179t

World Bank  
  GBD study (1990), 2–4, 35, 45, 50, 93, 399  
  GBD study (2001), 5–11, 35, 45–46  
  *World Development Report 1993: Investing in Health* (World Bank), 6, 427  
  *World Development Report 2003* (World Bank), 46  
World Health Organization (WHO)  
  on assessment of global burden of disease, 45  
  CHERG. *See* Child Health Epidemiology Reference Group  
  on HIV/AIDS mortality, 63  
  Multicountry Survey Study, 93  
  World Health Survey, 93  
  *World Health Report 2001*, 45  
  *World Health Report 2002*, 65, 246  
  *World Health Report 2003*, 45  
  
years of healthy life lost as result of disability (YLD)  
  burden of, 85–87  
  comparison of high-income to low- and middle-income countries, 87f, 92  
  data sources for, 75–79, 77–78t  
  disability weights for diseases and conditions, 119–125t  
  uncertainty in, 417, 420  
  
ensuring consistency by using DisMod, 74–75, 75f  
estimation of, 48, 73–74  
  by region (2001), 75  
incidence and prevalence, calculation of, 73–85  
leading causes of, 85–86, 86t  
neonatal mortality and stillbirths, 439–441, 447t  
regional variation in life expectancy and, 86–87, 87f  
years of life lost (YLL)  
  age weighting in, 401–402, 401f, 402f, 402t, 405f  
  defined, 48  
  discounting in, 401–402, 401f, 402t  
  estimation of, 48  
  neonatal mortality and stillbirths, 439–441, 440f, 440t, 446t, 449–451t, 455–457t  
  premature death and, 73, 73f  
  by region, 87–90, 87f  
risk factors and burden of disease, 269–393t  
  
zinc deficiency as risk factor, 287–290t

